



## Microalgal Polysaccharides: A Review of Chemical Structure, Biological Activities, and Prospects for Commercial Production

Aminikhoei.Z<sup>\*1</sup>, Mirbakhsh.M<sup>2</sup>, Pormozafar.S<sup>3</sup>, Khallagi.B<sup>4</sup>

1- Agricultural Research Education and Extension Organization (AREEO), Iranian Fisheries Science Research Institute (IFSRI), offshore fisheries Research Center, Chabahar, Iran.

2- Iranian Fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran.

3- Persian Gulf Mollusks Research Station, Persian Gulf and Oman Sea Ecology Research Center, Agricultural Research Education and Extension Organization (AREEO), Iranian Fisheries Sciences Research Institute (IFSRI), Bandar- e- Lengeh, Iran.

4- Makran Blue Algae Company, Science and Technology Park of Sistan and Baluchestan Province, Chabahar Campus, Iran.

### Abstract:

Microalgae have attracted significant attention from researchers in recent decades due to their rich bioactive compounds. Characteristics such as high growth rates, the ability to be cultivated in diverse environments, and the lack of direct competition with human food sources make them an attractive option for sustainable production. However, achieving economic and commercial viability for microalgal-derived products requires a thorough examination of biological, technological, and economic factors. This review explores the structural features, biological activities, and economic aspects of microalgal-derived polysaccharides. Polysaccharides extracted from microalgae exhibit considerable differentiation from land plant polysaccharides due to the heteropolymeric nature of their constituent monosaccharides and their high sulfate content. Moreover, numerous studies have shown that the diversity and specificity of the chemical structures of these compounds result in a range of bioactivities, including antibacterial, antiviral, and anticancer properties. Despite the significant biological advantages of these compounds, the large-scale and commercial production of microalgal polysaccharides faces technical and economic challenges that must be addressed.

### Review History:

Received: 9/16/2025

Revised: 10/12/2025

Accepted: 10/26/2025

Available Online: 10/28/2025

### Keywords:

Microalgae  
Polysaccharides  
Chemical Structure  
Biological Activities  
Economic Production

### How To Cite This Article:

Aminikhoei, Z., Mirbakhsh, M., Pormozafar, S., & Khallagi, B. (2025). Microalgal Polysaccharides: A Review of Chemical Structure, Biological Activities, and Prospects for Commercial Production. *Int. J. Biotech. Adv. Res.*, 1(1) 79-85



\*Corresponding author's email: zamini.41@gmail.com



Copyrights for this article are retained by the author(s) with publishing rights granted to Green Wave Pub. The content of this article is subject to the terms and conditions of the Creative Commons Attribution 4.0 International (CC BY 4.0) License. For more information, please visit <https://creativecommons.org/licenses/by/4.0/>

### 1.1. Review of the biological potential of marine microalgae and their competitive advantages relative to terrestrial plants

The exponential growth of the global population, the increasing demand for food security and public health, and the emergence of drug resistance and environmental pollution have exerted substantial pressure on traditional sources of bioactive compounds. Although terrestrial plant resources offer certain advantages, their cultivation is constrained by fundamental challenges such as the need for arable land, high freshwater consumption, and seasonal and climatic effects on the quantity and quality of produced compounds. These limitations have necessitated exploration of alternative sources, particularly aquatic environments. The oceans, which cover more than 70% of the Earth's surface, represent a vast repository of largely untapped biodiversity, encompassing an estimated ~800,000 species of microalgae and cyanobacteria (Kumar et al, 2020; Ravindran et al., 2021; Silvello et al., 2022). Microalgae, as efficient biological factories, are capable of producing a unique suite of metabolites including polysaccharides, pigments, unsaturated fatty acids, and proteins and therefore constitute a valuable resource for pharmaceutical, food, cosmetic, biofuel, and bioremediation applications. The global microalgae market was estimated at approximately USD 3.4 billion in 2020 and is projected to reach USD 4.6 billion by 2027, a growth attributed to increasing awareness of the environmental benefits and economic applications of microalgae (Sasaki et al, 2020; Koçer et al., 2021; Li et al., 2022).

Microalgae, as unicellular photoautotrophic organisms, offer unparalleled potential to address environmental and industrial challenges. Among the bioactive metabolites produced by microalgae, their polysaccharide derivatives characterized by unique chemical structures and biological activities are of particular interest. The following

sections present a detailed examination of the structural diversity and biological properties of microalgal polysaccharides (Chamizo et al, 2020; Deamicci et al., 2021; Gouda et al., 2022).

### 2.1. Structural Characteristics of Microalgal Polysaccharides

Polysaccharides are polymers composed of long chains of monosaccharides and are classified into homopolysaccharides (composed of a single type of monosaccharide) and heteropolysaccharides (containing two or more types of monomeric units). The structure and composition of the polysaccharide are determined by the types of monosaccharides present in the chain and the nature of the linkages between them. Microalgal polysaccharides are primarily heteropolymers consisting of galactose, xylose, and glucose in varying ratios, which are connected through glycosidic bonds. Other sugars, such as rhamnose, fucose, arabinose, mannose, orthomethylated sugars, and acidic residues of glucuronic acid and galacturonic acid, may also be components of the polysaccharides found in microalgae and cyanobacteria. Glucose is the most commonly encountered sugar. Fructose is generally not found in microalgal exopolysaccharides but is often a component of exopolysaccharides produced by cyanobacteria (Morales-Jiménez et al., 2020; Rachidi et al., 2020; Morais et al., 2020; Morais et al., 2021). Polysaccharides extracted from microalgae exhibit significant structural and molecular differences from those derived from other sources such as plants and animals. These polysaccharides often have a highly heterogeneous structure, meaning that the types of monosaccharides, the degree of branching, and the nature of glycosidic linkages can vary, exhibiting greater diversity than plant polysaccharides like starch or cellulose. A notable feature of microalgal polysaccharides is the presence of sulfate groups in their structure, which are rarely found in plant or animal sources.

Microalgal polysaccharides typically have variable molecular weights and specific branching patterns. The structural features of microalgal polysaccharides are influenced by the taxonomy of the algae and the cultivation conditions. Overall, these structural differences make microalgal polysaccharides exhibit unique biological and pharmaceutical functions compared to plant or animal polysaccharides. Microalgal polysaccharides can be categorized into intracellular polysaccharides and extracellular polysaccharides (Medina-Cabrera et al., 2020)

### 3.1. Bioactivities of Microalgal Polysaccharides and Their Relationship with Chemical Structure

Polysaccharides are natural macromolecules that participate in numerous biological processes such as cell adhesion, cell-to-cell communication, and the immune response. Furthermore, these bioactive compounds possess antioxidant, immunomodulatory, anticancer, antimicrobial, anti-inflammatory, and antiviral properties. Microalgal polysaccharides can be intracellular or extracellular (Exopolysaccharides EPS). Microalgal EPS are typically heteropolysaccharides with a high diversity of sugars (including xylose, rhamnose, and galactose) and often contain sulfated groups and uronic acids. The chemical characteristics of microalgal polysaccharides have a significant effect on their biological activity. The functional properties of polysaccharides vary based on their monosaccharide composition, molecular weight, sulfate and uronic acid content, linkage types, and distribution within the molecule (Li et al., 2020). The presence of rare monosaccharide constituents and oligosaccharides in microalgae, such as rhamnose or fucose, often confers biological activities. The presence of sulfate groups or acidic monosaccharides (such as uronic acids) also has a significant effect on their biological activity. For example, the increased net

charge of the EPS from the alga *Porphyridium cruentum* (resulting from sulfate and acidic monosaccharides) has led to enhanced antitumor and antiviral activities (Levasseur et al., 2020; Prybylski et al., 2020; Kazachenko et al., 2021; Yi et al., 2021).

In addition, chain size, spatial conformation, and the presence of other chemical groups (e.g., amino acids, proteins, or nucleic acids) are factors influencing their properties. For example, the antioxidant potential of the *Porphyridium cruentum* exopolysaccharide is associated with its sulfate content (4.5%) and/or the presence of a 66 kDa glycoprotein. Furthermore, the antioxidant property of polysaccharides can be inversely proportional to their molecular weight; a study revealed that the 6.5 kDa fragment exhibited higher radical scavenging activity than the 256 kDa and 60.6 kDa fragments. In a study, the polysaccharide from *Porphyridium* sp. exhibited antibacterial activity against *Escherichia coli* (72%) and *Bacillus subtilis* (35% reduction). The authors attributed this capacity to the different cell wall composition and structure of the bacteria. Regarding the antiviral potential of microalgal polysaccharides, strains with higher sulfation and uronic acid levels showed greater activity. Uronic acid constituents, sulfate half-ester groups, and carboxyl groups confer anionic properties to exopolysaccharides, ultimately enabling them to act as protective agents against viruses (Costa et al., 2020; Alvarez et al., 2021; Madadi et al., 2021).

### 4.1. Prospects for Economic Production of Polysaccharides from Microalgae

According to various reports, the global market size for algae (including microalgae and macroalgae) in 2022 was estimated at approximately 1.5 to 2 billion dollars. The compound annual growth rate for this market is projected to be between 6% and 10% for the period 2023 to 2030 or 2023 to 2032. This growth rate indicates a positive

outlook, driven by the increasing demand for natural, sustainable, and environmentally friendly products. However, the path to commercialization is confronted with numerous challenges. The first hurdle is the high cost of industrial-scale production, ranging from cultivation and harvesting to extraction and purification. A second challenge is the competition from cheaper alternative sources, such as vegetable oils or synthetic compounds. Furthermore, stringent regulations in the food and pharmaceutical safety sectors can impede the market entry process for new products (Guo et al., 2021; Colusse et al., 2021).

In the industrial-scale production of the primary feedstock, the type of cultivation system and production efficiency play a decisive role. Temperature and oxygen management, equipment sanitation, and biomass separation from the culture medium are among the key technical challenges in this industry. The choice between open systems (such as raceway ponds) and closed systems (tubular or flat-panel photobioreactors) directly impacts cost and efficiency. Open ponds, owing to their lower capital investment and ease of operation, are suitable for the mass production of low-value-added biomass (such as animal feed or biofertilizers); however, controlling cultivation conditions is difficult, and they are prone to contamination and fluctuations in product quality. In contrast, photobioreactors enable more precise control over growth conditions, increased biomass density, and the production of high-value compounds such as bioactive polysaccharides, although their capital and energy costs are higher (Parwani et al., 2021). An appropriate system design must maximize light and carbon dioxide utilization while minimizing mass transfer limitations. Furthermore, the scalability of these systems is critical; many technologies that are successful at the laboratory scale face challenges such as efficiency losses and increased costs at the industrial scale. Therefore, the choice of a

cultivation system must be based on the type of target product, its economic value, and regional conditions (access to land, water, and energy). The optimal balance among capital investment, production efficiency, and product quality is the key to achieving economically viable and sustainable microalgal production (Kusmayadi et al., 2021).

In terms of the technologies required for industrial production, the downstream processes of harvesting, drying, and extraction are determinants of the final cost following biomass production. Microalgal harvesting is considered the most cost-intensive step due to the minute size of the cells and low biomass concentration. Drying also impacts quality and cost; freeze-drying preserves the quality of sensitive compounds but is costly, whereas spray-drying is more economical but is accompanied by the risk of degrading bioactive metabolites. In the extraction stage, the use of green solvents such as ethanol or supercritical carbon dioxide, and novel technologies (ultrasound, microwave), can enhance process efficiency and sustainability. The biorefinery approach, in which multiple products (proteins, lipids, pigments) are extracted from a single biomass, is the primary key to reducing costs and increasing value addition. Conversely, significant economic opportunities also exist. The development of novel harvesting and extraction technologies, the application of biorefinery systems for the co-production of multiple products, and the utilization of wastewaters or low-cost carbon sources can reduce costs. Furthermore, increasing consumer awareness regarding health and environmental sustainability will bolster the market for these products. In summary, the economic future for producing bioactive compounds from microalgae is promising, but its realization will require technological innovation, supportive policies, and sustainable business models (Costa et al., 2021).

A precise understanding of the target market, market size, competitive pricing, and the needs of end customers is essential for determining product profitability. Does the product in question have a position in the current market, or does it necessitate the creation of a new one? Determining the product's value-added and its differentiation from competing products is also important. The greatest economic challenge arises

when microalgal products must compete with similar products produced from conventional or chemical sources. The utilization of microalgae as a source for specific active compounds is often not economically justifiable in comparison to conventional or synthetic alternatives, owing to the high costs of biomass production and purification (Moreira et al., 2022).

**Table 1: Commercial production of microalgal pigments**

Production Capacity (ton/year)	Year Established	Location/ Country	Company Name	Product	Microalgae
n.a.	1990	Kailua-Kona, Hawaii	Nutrex Hawaii	Hawaiian <i>Spirulina</i> , BioAstin Hawaiian Astaxanthin	<i>Spirulina</i> , <i>Haematococcus pluvialis</i>
18	2003	China	Jingzhou Natural Astaxanthin Inc.	Astaxanthin powder (1.5 to 3.0%), natural astaxanthin oleoresin, natural astaxanthin softgels (5 to 10%)	<i>Haematococcus pluvialis</i>
n.a.	n.a.	n.a.	Algatech	n.a.	<i>Haematococcus pluvialis</i> , <i>Phaeodactylum tricornutum</i> , <i>Porphyridium cruentum</i> , <i>Nannochloropsis</i> sp.
n.a.	1969	Osaka, Japan	Sun <i>Chlorella</i>	<i>Chlorella</i> tablets and drinks	<i>Chlorella</i>
<i>Chlorella</i> 1000, <i>Spirulina</i> 200	1967	Taiwan	Far East Microalgae Ind. Co., Ltd.	Organic <i>Spirulina</i> and <i>Chlorella</i> tablets, food supplements, and aquaculture feed	<i>Chlorella</i> and <i>Spirulina</i>
0.9	1995	China	Beijing Gingko Group	Astaxanthin	<i>Haematococcus pluvialis</i>
1.2	n.a.	China	Stone Forest Astaxanthin Biotech Co., Ltd.	Astaxanthin	<i>Haematococcus pluvialis</i>
0.6	n.a.	China	Yunnan Alphy Biotech Co., Ltd.	Astaxanthin	<i>Haematococcus pluvialis</i>
0.4	n.a.	China	Yunnan SGYJ Biotech Co., Ltd	Astaxanthin	<i>Haematococcus pluvialis</i>

**Table 2.** Market size and potential market for microalgal products (Patel et al., 2022)

Market Size (2009) (Million US\$)	Potential Market (2020) (Million US\$)	Sectors/Market Fields
300	800	Colorants
30	300	Nutraceuticals
Developing	500	Pharmaceuticals/Chemicals
Emerging/rising	30	Cosmetics/Personal care

Currently, markets are segmented into three categories: 1. High-value products such as pigments and supplements; these are profitable and suitable for current investment. 2. Food and chemical products, for which competitiveness requires that production costs be lowered, a goal achievable within the next decade. 3. Biofuels; this sector is not currently profitable and requires further progress in cost reduction (Patel et al., 2022).

## References

- Alvarez, A. L., Weyers, S. L., Goemann, H. M., Peyton, B. M., & Gardner, R. D. (2021). Microalgae, soil and plants: A critical review of microalgae as renewable resources for agriculture. *Algal Research*, 54, 102200.
- Chamizo, S., Adessi, A., Torzillo, G., & De Philippis, R. (2020). Exopolysaccharide features influence growth success in bio crust-forming cyanobacteria, moving from liquid culture to sand microcosms. *Frontiers in Microbiology*, 11, 568224.
- Colusse, G. A., Carneiro, J., Duarte, M. E. R., De Carvalho, J. C., & Nosedá, M. D. (2021). Advances in microalgal cell wall polysaccharides: A review focused on structure, production, and biological application. *Critical Reviews in Biotechnology*, 41(1), 1–16.
- Costa, J. A. V., Lucas, B. F., Alvarenga, A. G. P., Moreira, J. B., & Morais, M. G. (2021). Microalgae polysaccharides: An overview of production, characterization, and potential applications. *Polysaccharides*, 2(4), 759–772.
- Costa, J. A. V., Moreira, J. B., Fanka, L. S., Kosinski, R. D. C., & de Morais, M. G. (2020). Microalgal biotechnology applied in biomedicine. In O. Konur (Ed.), *Handbook of algal science, technology and medicine* (pp. 429–439). Elsevier.
- Deamici, K., de Morais, M., Santos, L., Muylaert, K., Gardarin, C., Costa, J., & Laroche, C. (2021). Static magnetic fields effects on polysaccharides production by different microalgae strains. *Applied Sciences*, 11(12), 5299.
- Gouda, M., Tadda, M. A., Zhao, Y., Farmanullah, F., Chu, B., Li, X., & He, Y. (2022). Microalgae bioactive carbohydrates as a novel sustainable and eco-friendly source of prebiotics: Emerging health functionality and recent technologies for extraction and detection. *Frontiers in Nutrition*, 9, 806692.
- Guo, W., Zhu, S., Li, S., Feng, Y., Wu, H., & Zeng, M. (2021). Microalgae polysaccharides ameliorates obesity in association with modulation of lipid metabolism and gut microbiota in high-fat-diet fed C57BL/6 mice. *International Journal of Biological Macromolecules*, 182, 1371–1383.
- Kazachenko, A. S., Akman, F., Malyar, Y. N., Issaoui, N., Vasilieva, N. Y., & Karacharov, A. A. (2021). Synthesis optimization, DFT and physicochemical study of chitosan sulfates. *Journal of Molecular Structure*, 1245, 131083.
- Koçer, A. T., Inan, B., Usul, S. K., Özçimen, D., Yılmaz, M. T., & Işıldak, I. (2021). Exopolysaccharides from microalgae: Production, characterization, optimization and techno-economic assessment. *Brazilian Journal of Microbiology*, 52(4), 1779–1790.
- Kumar, M., Sun, Y., Rathour, R., Pandey, A., Thakur, I. S., & Tsang, D. C. W. (2020). Algae as potential feedstock for the production of biofuels and value-added products: Opportunities and challenges. *Science of The Total Environment*, 716, 137116.
- Kusmayadi, A., Leong, Y. K., Yen, H.-W., Huang, C.-Y., & Chang, J.-S. (2021). Microalgae as sustainable food and feed sources for animals and humans: Biotechnological and environmental aspects. *Chemosphere*, 271, 129800.
- Levasseur, W., Perré, P., & Pozzobon, V. (2020). A review of high value-added molecules production by microalgae in light of the classification. *Biotechnology Advances*, 41, 107545.

14. Li, S., Ji, L., Chen, C., Zhao, S., Sun, M., Gao, S., Gao, Z., Wu, H., & Fan, J. (2022). Efficient accumulation of high-value bioactive substances by carbon to nitrogen ratio regulation in marine microalgae *Porphyridium purpureum*. *Bioresource Technology*, 347, 126620.
15. Li, T. T., Huang, Z. R., Jia, R. B., Lv, X. C., Zhao, C., & Liu, B. (2021). *Spirulina platensis* polysaccharides attenuate lipid and carbohydrate metabolism disorder in high-sucrose and high-fat diet-fed rats in association with intestinal microbiota. *Food Research International*, 147, 110530.
16. Li, Y., Wang, C., Liu, H., Su, J., Lan, C. Q., Zhong, M., & Hu, X. (2020). Production, isolation and bioactive estimation of extracellular polysaccharides of green microalga *Neochloris oleoabundans*. *Algal Research*, 48, 101883.
17. Madadi, R., Maljaee, H., Serafim, L. S., & Ventura, S. P. M. (2021). Microalgae as contributors to produce biopolymers. *Marine Drugs*, 19(8), 466.
18. Medina-Cabrera, E. V., Rühmann, B., Schmid, J., & Sieber, V. (2020). Optimization of growth and EPS production in two *Porphyridium* strains. *Bioresource Technology Reports*, 11, 100486.
19. Morais, M. G., Morais, E. G., Cardias, B. C., Vaz, B. S., Moreira, J. B., Mitchell, G., & Costa, J. A. V. (2020). Microalgae as a source of sustainable biofuels. In V. K. Gupta, H. Treichel, & S. Rodriguez-Couto (Eds.), *Recent developments in bioenergy research* (pp. 253–271). Elsevier.
20. Morais, M. G., Rosa, G. M., Moraes, L., Alvarenga, A. G. P., Silva, J. L. V., & Costa, J. A. V. (2021). Microalgae polysaccharides with potential biomedical application. In *Polysaccharides of microbial origin* (pp. 363–380). Springer International Publishing.
21. Morales-Jiménez, M., Gouveia, L., Yáñez-Fernández, J., Castro-Muñoz, R., & Barragán-Huerta, B. E. (2020). Production, preparation and characterization of microalgae-based biopolymer as a potential bioactive film. *Coatings*, 10(2), 120.
22. Moreira, J. B., Kuntzler, S. G., Bezerra, P. Q. M., Cassuriaga, A. P. A., Zapparoli, M., da Silva, J. L. V., Costa, J. A. V., & da Morais, M. G. (2022). Recent advances of microalgae exopolysaccharides for application as bioflocculants. *Polysaccharides*, 3(1), 264–276.
23. Parwani, L., Bhatt, M., & Singh, J. (2021). Potential biotechnological applications of cyanobacterial exopolysaccharides. *Brazilian Archives of Biology and Technology*, 64, e21200401.
24. Patel, A. K., Vadrade, A. P., Singhanian, R. R., Michaud, P., Pandey, A., Chen, S. J., Chen, C. W., & Dong, C. D. (2022). Algal polysaccharides: Current status and future prospects. *Phytochemistry Reviews*, 21(2), 451–483.
25. Prybylski, N., Toucheteau, C., El Alaoui, H., Bridiau, N., Maugard, T., Abdelkafi, S., Fendri, I., Delattre, C., Dubessay, P., & Pierre, G. (2020). Bioactive polysaccharides from microalgae. In E. Jacob-Lopes, L. Q. Zepka & M. I. Queiroz (Eds.), *Handbook of microalgae-based processes and products* (pp. 533–571). Academic Press.
26. Ravindran, R., & Rajauria, G. (2021). Carbohydrates derived from microalgae in the food industry. In C. Galanakis (Ed.), *Cultivated microalgae for the food industry* (pp. 127–146). Academic Press.
27. Sasaki, M., Takagi, A., Ota, S., Kawano, S., Sasaki, D., & Asayama, M. (2020). Coproduction of lipids and extracellular polysaccharides from the novel green alga *Parachlorella* sp. BX1.5 depending on cultivation conditions. *Biotechnology Reports*, 25, e00392.
28. Silvello, M. A. D. C., Gonçalves, I. S., Azambuja, S. P. H., Costa, S. S., Silva, P. G. P., Santos, L. O., & Goldbeck, R. (2022). Microalgae-based carbohydrates: A green innovative source of bioenergy. *Bioresource Technology*, 344, 126304.
29. Yi, Z., Su, Y., Brynjolfsson, S., Olafsdóttir, K., & Fu, W. (2021). Bioactive polysaccharides and their derivatives from microalgae: Biosynthesis, applications, and challenges. *Studies in Natural Products Chemistry*, 71, 67–8